

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

AMENDMENTS TO THE CLAIMS


This listing of claims will replace all prior versions, and listings, of claims in the application:

Listing of Claims:

1. (Cancelled)

3. - 6. (Cancelled)

7. (Currently amended) A method for the treatment of a disorder of the central nervous system related to or affected by the 5-HT₆ receptor wherein said disorder is selected from the group consisting essentially of: schizophrenia and depression; in a patient in need thereof which comprises administering to said patient a therapeutically effective amount of a compound of formula I.



(I)

wherein

A is N;

X is CR₁₁ or N;

Y is CR₇ or N with the proviso that when X is N, then Y must be CR₇;

R₁ is H, C₁-C₆alkylcarbonyl, C₁-C₆alkylcarbonyloxy or an C₁-C₆alkyl, C₁-C₆alkenyl, C₁-C₆alkynyl or cycloheteroalkyl group each optionally substituted;

R₂, R₃, R₄, R₅ and R₆ are each independently H, halogen, OH or an optionally substituted C₁-C₆alkyl group;

~~R₇ and R₁₁ are each independently~~ is H, halogen or an C₁-C₆alkyl, aryl, heteroaryl or C₁-C₆alkoxy group each optionally substituted;

R₈ is an C₁-C₆alkyl, aryl or heteroaryl group each optionally substituted;

R₉ is H, halogen or an C₁-C₆alkyl, C₁-C₆alkoxy, C₁-C₆alkenyl, aryl or heteroaryl group each optionally substituted;

R₁₀ is H, OH or an optionally substituted C₁-C₆alkoxy group;

m is an integer of 2;

n is 0 or an integer of 1, 2 or 3; and


--- represents a single bond or a double bond; or a pharmaceutically acceptable salt thereof.

12. (Cancelled)

14. - 17. (Cancelled)

18. (Currently Amended) A method for the preparation of a compound of formula

I.



wherein

A is N;

X is CR₁₁ or N;

Y is CR₇ or N with the proviso that when X is N, then Y must be CR₇;

R₁ is C₁-C₆alkylcarbonyl, C₁-C₆alkylcarbonyloxy or an C₁-C₆alkyl, C₁-C₆alkenyl, C₁-C₆alkynyl or cycloheteroalkyl group each optionally substituted;

R₂, R₃, R₄, R₅ and R₆ are each independently H, halogen, OH or an optionally substituted C₁-C₆alkyl group;

~~R₇ and R₁₁ are each independently~~ is H, halogen or an C₁-C₆alkyl, aryl, heteroaryl or alkoxy group each optionally substituted;

R₈ is an C₁-C₆alkyl, aryl or heteroaryl group each optionally substituted;

R_9 is H, halogen or an C_1 - C_6 alkyl, C_1 - C_6 alkoxy, C_1 - C_6 alkenyl, aryl or heteroaryl group each optionally substituted;


R_{10} is H, OH or an optionally substituted C_1 - C_6 alkoxy group;

m is an integer of 2;

n is 0 or an integer of 1, 2 or 3; and


--- represents a single bond ~~or a double bond~~

said method which comprises reacting a compound of formula Ia



wherein A, X, R_2 , R_3 , R_4 , R_5 , R_6 , R_7 , R_8 , R_9 , m and n are as defined hereinabove for formula I with a compound R_1 -Hal wherein R_1 is as defined hereinabove for formula I and Hal is Cl, Br or I.

19. (Currently Amended) A compound of formula I



wherein

A is N;

X is CR_{11} ;

Y is N;

R₁ is H, C₁-C₆alkylcarbonyl, C₁-C₆alkylcarbonyloxy or an C₁-C₆alkyl, C₁-C₆alkenyl, C₁-C₆alkynyl or cycloheteroalkyl group each optionally substituted;

R₂, R₃, R₄, R₅ and R₆ are each independently H, halogen, OH or an optionally substituted C₁-C₆alkyl group;

R₁₁ is H, halogen or an C₁-C₆alkyl, aryl, heteroaryl or C₁-C₆alkoxy group each optionally substituted;

R₈ is an C₁-C₆alkyl, aryl or heteroaryl group each optionally substituted;

R₉ is H, halogen or an C₁-C₆alkyl, C₁-C₆alkoxy, C₁-C₆alkenyl, aryl or heteroaryl group each optionally substituted;

R₁₀ is H, OH or an optionally substituted C₁-C₆alkoxy group;

m is an integer of 2;

n is 0 or an integer of 1, 2 or 3; and

--- represents a single bond ~~or a double bond~~; or
a pharmaceutically acceptable salt thereof.

20. (Previously presented) The compound according to claim 19 wherein R₈ is an optionally substituted phenyl group.

21. (Previously presented) The compound according to claim 19 selected from the group consisting of:

1-(phenylsulfonyl)-5-piperazin-1-yl-1H-indazole;

1-(phenylsulfonyl)-6-piperazin-1-yl-1H-indazole;

1-[(2-bromophenyl)sulfonyl]-6-piperazin-1-yl-1H-indazole;

1-[(4-bromophenyl)sulfonyl]-5-piperazin-1-yl-1H-indazole;

1-[(4-bromophenyl)sulfonyl]-6-piperazin-1-yl-1H-indazole;

1-[(5-bromothien-2-yl)sulfonyl]-5-piperazin-1-yl-1H-indazole;

1-[(5-bromothien-2-yl)sulfonyl]-6-piperazin-1-yl-1H-indazole;

1-[(4-fluorophenyl)sulfonyl]-5-piperazin-1-yl-1H-indazole;

1-[(4-fluorophenyl)sulfonyl]-6-piperazin-1-yl-1H-indazole;

methyl 4-[(5-piperazin-1-yl-1H-indazol-1-yl)sulfonyl]phenyl ether;


1-phenylsulfonyl-4-(4-propylpiperazin-1-yl)-1H-indazole;

1-phenylsulfonyl-4-piperazin-1-yl-1H-indazole;

1-phenylsulfonyl-4-(4-phenethylpiperazin-1-yl)-1H-indazole;

1-phenylsulfonyl-4-[4-(3-phenylpropyl)-piperazin-1-yl]-1H-indazole; and
the pharmaceutically acceptable salts thereof.

22. (Currently Amended) A pharmaceutical composition which comprises a
pharmaceutically acceptable carrier and an effective amount of a compound of formula I.



wherein

A is N;

X is CR₁₁;

Y is N;

R₁ is H, C₁-C₆alkylcarbonyl, C₁-C₆alkylcarbonyloxy or an C₁-C₆alkyl, C₁-C₆alkenyl, C₁-C₆alkynyl or cycloheteroalkyl group each optionally substituted;

R₂, R₃, R₄, R₅ and R₆ are each independently H, halogen, OH or an optionally substituted C₁-C₆alkyl group;

R₁₁ is H, halogen or an C₁-C₆alkyl, aryl, heteroaryl or C₁-C₆alkoxy group each optionally substituted;

R₈ is an C₁-C₆alkyl, aryl or heteroaryl group each optionally substituted;

R₉ is H, halogen or an C₁-C₆alkyl, C₁-C₆alkoxy, C₁-C₆alkenyl, aryl or heteroaryl group each optionally substituted;

R₁₀ is H, OH or an optionally substituted C₁-C₆alkoxy group;

m is an integer of 2;

n is 0 or an integer of 1, 2 or 3; and

--- represents a single bond or a double bond; or

a pharmaceutically acceptable salt thereof.

23. (Previously presented) The composition according to claim 22 having a
compound of formula I selected from the group consisting of:

1-(phenylsulfonyl)-5-piperazin-1-yl-1H-indazole;
1-(phenylsulfonyl)-6-piperazin-1-yl-1H-indazole;
1-[(2-bromophenyl)sulfonyl]-6-piperazin-1-yl-1H-indazole;
1-[(4-bromophenyl)sulfonyl]-5-piperazin-1-yl-1H-indazole;
1-[(4-bromophenyl)sulfonyl]-6-piperazin-1-yl-1H-indazole;
1-[(5-bromothien-2-yl)sulfonyl]-5-piperazin-1-yl-1H-indazole;
1-[(5-bromothien-2-yl)sulfonyl]-6-piperazin-1-yl-1H-indazole;
1-[(4-fluorophenyl)sulfonyl]-5-piperazin-1-yl-1H-indazole;
1-[(4-fluorophenyl)sulfonyl]-6-piperazin-1-yl-1H-indazole;
methyl 4-[(5-piperazin-1-yl-1H-indazol-1-yl)sulfonyl]phenyl ether;
1-phenylsulfonyl-4-(4-propylpiperazin-1-yl)-1H-indazole;
1-phenylsulfonyl-4-piperazin-1-yl-1H-indazole;
1-phenylsulfonyl-4-(4-phenethylpiperazin-1-yl)-1H-indazole;
1-phenylsulfonyl-4-[4-(3-phenylpropyl)-piperazin-1-yl]-1H-indazole; and
the pharmaceutically acceptable salts thereof.